

ELEMENTARY SPECTRAL CALCULATIONS IN RINGS

GYULA LAKOS

ABSTRACT. We examine the validity of certain spectral integrals in topological rings. We consider the sign and square-root functions in polymetric rings containing $\frac{1}{2}$. It turns out that formal analogues of classical transformation kernels and the resolvent identity can be used to understand the situation. In the lack of $\frac{1}{2}$ the functions $\frac{1}{2} - \frac{1}{2} \operatorname{sgn}(\frac{1}{2} - z)$ and $\frac{1}{2} - \sqrt{\frac{1}{4} - z}$ can be generalized, respectively.

0. INTRODUCTION

Spectral integrals like

$$(1) \quad \operatorname{sgn} Q = \int_{\{z \in \mathbb{C} : |z|=1\}} \frac{\frac{1-z}{2} + \frac{1+z}{2} Q}{\frac{1+z}{2} + \frac{1-z}{2} Q} \frac{|dz|}{2\pi}$$

and

$$(2) \quad \sqrt{S} = \int_{\{z \in \mathbb{C} : |z|=1\}} \frac{S}{\frac{1}{z} \left(\left(\frac{1+z}{2} \right)^2 - \left(\frac{1-z}{2} \right)^2 S \right)} \frac{|dz|}{2\pi}$$

are often useful. They extend the complex functions $\operatorname{sgn} Q$, which is the sign function of (the real part of) Q , and $\sqrt{S}$, which is the square root function cut along the negative real axis. Definitions like above are justified if they are supported by appropriate algebraic identities and spectral properties. This is the situation in linear analysis where the formulas above can be established for elements with appropriate spectral properties in great generality, even if the resolvent terms are not necessarily continuous, cf. Haase [1], Martínez Carracedo–Sanz Alix [3]. We refer to this case as the “analytic case”.

However, in the analytic case, if the resolvent terms are continuous then they are also smooth and one can expand everything in terms of Fourier series, or rather Laurent series in z . One can naturally ask if similar computations can be done in more general rings, in particular, in rings without a natural $\mathbb{R}$ -action. It is natural to check these ideas against formal Laurent series on polymetric rings. We refer to this case as the “algebraic case”.

We call a ring $\mathfrak{A}$ polymetric if

- a.) its topology is induced by a family of “seminorms” $p : \mathfrak{A} \rightarrow [0, +\infty)$ such that $p(0) = 0$, $p(-X) = p(X)$, $p(X + Y) \leq p(X) + p(Y)$,
 - b.) for each “seminorm” p there exists a “seminorm” $\tilde{p}$ such that $p(XY) \leq \tilde{p}(X)\tilde{p}(Y)$;
- ie. it is a polymetric space whose multiplication is compatible with the topology. Dealing with Laurent series in z integration over the unit circle becomes a formal process, it is nothing else but detecting the coefficient of z^0 . On the other hand, to the multiplication of Laurent series some sort of convergence control is required. Primarily, we will be interested Laurent series with rapidly decreasing coefficients. A sequence is rapidly decreasing if it is rapidly decreasing in each seminorm. Furthermore, we consider only sequentially complete, Hausdorff polymetric rings. We also assume that $\frac{1}{2} \in \mathfrak{A}$. Then (1) and (2) are meaningful.

2000 *Mathematics Subject Classification*. Primary: 46H30, Secondary: 13J99.

Key words and phrases. Sign operation, square root operation, formal transformation kernels.

Indeed, applied to elements with appropriate “spectral” properties the expressions $\operatorname{sgn} Q$ and $\sqrt{S}$ will have good properties justifying the notation. This can be proved by an analysis of the coefficients, The algebraic approach is particularly manageable in the case of (†): it is essentially shown in Karoubi [2] by a direct analysis of coefficients that the expression $\operatorname{sgn} Q$ yields an involution compatible to factorization of affine loops. Nevertheless, such computations are not necessarily very enlightening. The objective of this paper is to prove our statements regarding the algebraic case and to do this in a manner which brings the algebraic and analytic cases together, at least formally. It turns out that the basic tool of the analytic case, the resolvent identity works generally. Another natural question is what happens in the lack of $\frac{1}{2}$. Then the sign and square root functions are not really appropriate. Instead, we can generalize

$$(3) \quad \operatorname{idem} P = \int_{\{z \in \mathbb{C} : |z|=1\}} \frac{Pz}{1 - P + Pz} \frac{|dz|}{2\pi}$$

and

$$(4) \quad \sqrt[T]{T} = \int_{\{z \in \mathbb{C} : |z|=1\}} \frac{(1+z)T}{1 + (z - 2 + z^{-1})T} \frac{|dz|}{2\pi},$$

extending the functions $\operatorname{idem} P = \frac{1}{2} - \frac{1}{2} \operatorname{sgn} \left(\frac{1}{2} - P \right)$ and $\sqrt[T]{T} = \frac{1}{2} - \sqrt{\frac{1}{4} - T}$.

1. LAURENT SERIES

1.1. If $\mathfrak{A}$ is a polymetric ring then we may consider formal Laurent series $a = \sum_{n \in \mathbb{N}} a_n z^n$. If p is a seminorm on $\mathfrak{A}$, and α is a non-negative function then we may define $p_\alpha(a) = \sum_{n \in \mathbb{N}} \alpha(n) p(a_n)$. Then we may consider the polymetric spaces a.) $\mathfrak{A}[z^{-1}, z]^f$ of essentially finite Laurent series, b.) $\mathfrak{A}[z^{-1}, z]^\infty$ of rapidly decreasing Laurent series, c.) $\mathfrak{A}[z^{-1}, z]^b$ of “summable” Laurent series, d.) $\mathfrak{A}[[z^{-1}, z]]^b$ of bounded Laurent series, e.) $\mathfrak{A}[[z^{-1}, z]]^\infty$ of polynomially growing Laurent series, f.) $\mathfrak{A}[[z^{-1}, z]]^f$ of formal Laurent series, as the spaces which contain series bounded for each p_α such that a.) α is unrestricted, b.) α is polynomially growing, c.) α is bounded, d.) α is summable, e.) α is rapidly decreasing, f.) α is vanishing except at finitely many places. We have continuous inclusions

$$\mathfrak{A}[z^{-1}, z]^f \hookrightarrow \mathfrak{A}[z^{-1}, z]^\infty \hookrightarrow \mathfrak{A}[z^{-1}, z]^b \hookrightarrow \mathfrak{A}[[z^{-1}, z]]^b \hookrightarrow \mathfrak{A}[[z^{-1}, z]]^\infty \hookrightarrow \mathfrak{A}[[z^{-1}, z]]^f.$$

Of these spaces $\mathfrak{A}[z^{-1}, z]^f$, $\mathfrak{A}[z^{-1}, z]^\infty$, $\mathfrak{A}[z^{-1}, z]^b$ will remain polymetric rings. Indeed, $\tilde{p}_\alpha$ can be chosen as $p_{\tilde{\alpha}}$, where $\tilde{\alpha}(n) = 1 \vee \max_{-n \leq m \leq n} |\alpha(m)|$. Nevertheless there are compatible pairings $\mathfrak{A}[z^{-1}, z]^\bullet \times \mathfrak{A}[[z^{-1}, z]]^\bullet \rightarrow \mathfrak{A}[[z^{-1}, z]]^\bullet$. Essentially the same applies to the spaces of power series $\mathfrak{A}[z]^f$, $\mathfrak{A}[z]^\infty$, $\mathfrak{A}[z]^b$, $\mathfrak{A}[[z]]^b$, $\mathfrak{A}[[z]]^\infty$, $\mathfrak{A}[[z]]^f$, except there even $\mathfrak{A}[[z]]^\infty$, $\mathfrak{A}[[z]]^f$ are polymetric rings. Indeed, for them $\tilde{p}_\alpha$ can be chosen as $p_{\tilde{\alpha}}$, where $\tilde{\alpha}(n) = \sqrt{\max_{n \leq m} |\alpha(m)|}$. An element like $\frac{1+Q}{2} + \frac{1-Q}{2}z$ may be considered either as an element of $\mathfrak{A}[z^{-1}, z]^f$, $\mathfrak{A}[z^{-1}, z]^\infty$, or $\mathfrak{A}[z^{-1}, z]^b$. Practically, the difference is that the larger the ring is the easier is to find a multiplicative inverse of the element given.

1.2. If $a(z) = \sum_{n \in \mathbb{Z}} a_n z^n \in \mathfrak{A}[[z^{-1}, z]]^f$ then we formally define

$$\int a(z) \frac{|dz|}{2\pi} = a_0.$$

The Hilbert kernel is defined as

$$\left[\frac{1+z}{1-z} \right] = \sum_{s \in \mathbb{Z}} (\operatorname{sgn} s) z^s.$$

1.3. We introduce some further terminology which may look like a bit grandiose at first sight, yet it is very useful. For $a(z) = \sum_{n \in \mathbb{Z}} a_n z^n \in \mathfrak{A}[z^{-1}, z]^b$ we let

$$\lim_{z \nearrow 1} a(z) = \sum_{n \in \mathbb{Z}} a_n.$$

Naturally, this notation also applies to power series.

In what follows we let $\mathfrak{A}[z^{-1}, z]$, $\mathfrak{A}[[z^{-1}, z]]$ denote $\mathfrak{A}[z^{-1}, z]^\infty$, $\mathfrak{A}[[z^{-1}, z]]^\infty$, respectively, but similar statements hold for f and b , too.

Proposition 1.4. For $a(t, z) \in \mathfrak{A}[t][[z^{-1}, z]]$ and $b(z) \in \mathfrak{A}[z^{-1}, z]$

$$\int \left(\lim_{t \nearrow 1} a(t, z) \right) b(z) \frac{|dz|}{2\pi} = \lim_{t \nearrow 1} \int (a(t, z)) b(z) \frac{|dz|}{2\pi}.$$

Proof. This is just the generalized associativity of the rapidly decreasing (hence absolute convergent) sum $\sum_{n \in \mathbb{N}, s \in \mathbb{Z}} a_{n,s} b_{-s}$. $\square$

1.5. For the sake of brevity we call elements of $\mathfrak{A}[t][[z^{-1}, z]]$ as transformation kernels. Practically, the convenient thing is to consider those elements of the ring $\mathfrak{A}[[t]]^f[z^{-1}, z]^b$ which can be thought to be transformation kernels. (This is advantageous from computational viewpoint, because the product of $a(t, z)$ and $b(z) \in \mathfrak{A}[z^{-1}, z]$ formally yields the same element of $\mathfrak{A}[[t]]^f[z^{-1}, z]^b$ either we interpret $a(t, z) \in \mathfrak{A}[[t]]^f[z^{-1}, z]^b$ or $a(t, z) \in \mathfrak{A}[t][[z^{-1}, z]]$, but the first case is often easier to compute with.)

Such elements are the Poisson kernel

$$\mathcal{P}(t, z) = \frac{1 - t^2}{(1 - tz)(1 - tz^{-1})} = \sum_{s \in \mathbb{Z}} t^{|s|} z^s,$$

the Hilbert-Poisson kernel

$$\mathcal{H}(t, z) = \frac{t(z - z^{-1})}{(1 - tz)(1 - tz^{-1})} = \sum_{s \in \mathbb{Z}} (\operatorname{sgn} s) t^{|s|} z^s,$$

the $\frac{1}{2}$ -shifted odd Poisson kernel

$$\mathcal{L}(t, z) = \frac{(1 - t)(1 + z)}{(1 - tz)(1 - tz^{-1})} = \sum_{s \in \mathbb{N}} t^s z^{-s} + \sum_{s \in \mathbb{N}} t^s z^{s+1},$$

and the variant regularization kernel

$$\tilde{\mathcal{R}}(t, z) = \frac{(1 + t)t(1 - z)(1 - z^{-1})}{2(1 - tz)(1 - tz^{-1})} = \sum_{s \in \mathbb{N}, s > 0} \frac{t^{s+1} - t^s}{2} z^{-s} + t + \sum_{s \in \mathbb{N}, s > 0} \frac{t^{s+1} - t^s}{2} z^s.$$

This latter one has the property

$$\lim_{t \nearrow 1} \tilde{\mathcal{R}}(t, z) = 1$$

(here we understand $\tilde{\mathcal{R}}(t, z) \in \mathfrak{A}[t][[z^{-1}, z]]$). The ordinary regularization kernel

$$\mathcal{R}(t, z) = \frac{t(1 - z)(1 - z^{-1})}{(1 - tz)(1 - tz^{-1})}$$

is just an element of $\mathfrak{A}[[t]]^f[z^{-1}, z]^b$, hence it is not so convenient algebraically. On the other hand, the use of variant regularization makes useful the variant Hilbert-Poisson kernel

$$\tilde{\mathcal{H}}(t, z) = \frac{1 + t}{2} \mathcal{H}(t, z) = \frac{(1 + t)t(z - z^{-1})}{2(1 - tz)(1 - tz^{-1})} = \sum_{s \in \mathbb{Z}} (\operatorname{sgn} s) \frac{1 + t}{2} t^{|s|} z^s.$$

It is not hard to see that the definitions and the proposition above can be formulated in the case when we have many variables $z, w, \dots$ instead of just z . For example we may consider the Hilbert kernel

$$\left[\frac{z+w}{z-w} \right] = \left[\frac{1+wz^{-1}}{1-wz^{-1}} \right] = \sum_{s \in \mathbb{Z}} (\text{sgn } s) w^s z^{-s}.$$

2. SPECTRAL CLASSES

2.1. In order to save some space we use the short-hand notation

$$\begin{aligned} \Lambda(a) &= \frac{1}{2}(1+a) \\ \Lambda(a, b) &= \frac{1}{2}(1+a+b-ab) \\ \Lambda(a, b, c) &= \frac{1}{4}(1+a+b+c-ab+ac-bc+abc) \\ \Lambda(a, b, c, d) &= \frac{1}{4}(1+a+b+c+d-ab-bc-cd+ac+ad+bd \\ &\quad + abc - acd - abd + bcd - abcd) \end{aligned}$$

etc., following the scheme

$$\Lambda(c_1, c_2, c_3, \dots, c_n) = 2^{-\lceil \frac{n}{2} \rceil} \sum_{\varepsilon \in \{0,1\}^n} \left(\prod_{1 \leq j < n} (-1)^{\varepsilon_j \varepsilon_{j+1}} \right) \left(\prod_{1 \leq k \leq n} c_k^{\varepsilon_k} \right)$$

such that the order of the symbols is preserved.

2.2. Let $\overline{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ denote the Riemann sphere. Some subsets are: $i\overline{\mathbb{R}} = i\mathbb{R} \cup \{\infty\}$, $\overline{\mathbb{R}}^- = (-\infty, 0] \cup \{\infty\}$, $\overline{\mathbb{C}}^- = \{s \in \mathbb{C} : \text{Re } s \leq 0\} \cup \{\infty\}$, $\mathring{\mathbb{D}}^1 = \{z \in \mathbb{C} : |z| < 1\}$. We define the functions $\text{pol } J = -i \text{sgn } iJ$, $|J|_i = \sqrt{-J^2}$, $|Q|_r = \sqrt{Q^2}$, $|P|_{\mathcal{F}} = \frac{1}{2} - |\frac{1}{2} - P|_r$. We have the following commutative diagram on certain subsets of the complex plane:

$$(5) \quad \begin{array}{ccccccc} & & & & J \in \mathbb{C} \setminus \overline{\mathbb{R}} & \xrightarrow{J \mapsto \text{pol } J} & J \in \{i, -i\} \\ & & & & \nearrow J \mapsto -J^2 & & \nearrow J \mapsto |J|_i \\ & & & & S \in \mathbb{C} \setminus \overline{\mathbb{R}}^- & \xrightarrow{S \mapsto \sqrt{S}} & Q \in \mathbb{C} \setminus \overline{\mathbb{C}}^- \\ & & & & \xleftarrow{Q \mapsto |Q|_r} & & \xleftarrow{Q \mapsto \text{sgn } Q} \\ & & & & Q \in \mathbb{C} \setminus i\overline{\mathbb{R}} & \xrightarrow{Q \mapsto \text{sgn } Q} & Q \in \{1, -1\} \\ & & & & \xleftarrow{Q \mapsto Q^2} & & \xleftarrow{Q \mapsto \frac{1-Q}{2}} \\ & & & & \xleftarrow{Q \mapsto \frac{1-Q}{2}} & & \xleftarrow{P \mapsto 1-2P} \\ & & & & T \in \mathbb{C} \setminus \left(\frac{1}{4} - \overline{\mathbb{R}}^-\right) & \xrightarrow{T \mapsto \sqrt[4]{T}} & P \in \mathbb{C} \setminus \left(\frac{1}{2} - \overline{\mathbb{C}}^-\right) \\ & & & & \xleftarrow{P \mapsto P(1-P)} & & \xleftarrow{P \mapsto |P|_{\mathcal{F}}} \\ & & & & P \in \mathbb{C} \setminus \left(\frac{1}{2} - i\overline{\mathbb{R}}\right) & \xrightarrow{P \mapsto \text{idem } P} & P \in \{0, 1\} \\ & & & & \xleftarrow{W \mapsto -W(1-W)^{-1}} & & \xleftarrow{P \mapsto -P(1-P)^{-1}} \\ & & & & W \in \mathring{\mathbb{D}}^1 & & \end{array}$$

such that $\text{pol } J$, $\text{sgn } Q$, $\text{idem } P$, $|J|_i$, $|Q|_r$, $|P|_{\mathcal{F}}$ yield idempotent operations and they yield decompositions

$$J = |J|_i \text{pol } J, \quad Q = |Q|_r \text{sgn } Q, \quad P = \text{idem } P + |P|_{\mathcal{F}} - 2|P|_{\mathcal{F}} \text{idem } P.$$

Definition 2.3. Suppose that $\mathfrak{A}$ is a locally convex algebra and R is a compact subset of $\overline{\mathbb{C}}$. We define $\text{Spec}_R(\mathfrak{A})$ as the set containing all elements $X \in \mathfrak{A}$ such that the functions

$$f_R : z \in R \setminus \{\infty\} \mapsto (z - X)^{-1} \quad \text{and} \quad g_R : z \in R \setminus \{0\} \mapsto X(1 - z^{-1}X)^{-1}$$

are well-defined and continuous. The topology of $\text{Spec}_R(\mathfrak{A})$ is induced from the compact-open topology of the continuous functions $(f_R)|_{\mathbb{D}^1 \cap R}$ and $(g_R)|_{R \setminus \mathring{\mathbb{D}}^1}$. (If R is symmetric for

conjugation then we may use the functions $f_R(z)f_R(\bar{z})$ and $g_R(z)g_R(\bar{z})$ to get a formally real characterization.)

Classes of interest are like $\widetilde{\text{Spec}}_{i\mathbb{R}}(\mathfrak{A})$, etc., ie. the elements spectrally avoiding $i\mathbb{R}$, etc. Another way to specify spectral conditions is to ask for skew-involutions, involutions, or idempotents. The main classes correspond to the sets in (5) for $\mathfrak{A} = \mathbb{C}$.

Definition 2.4. We define the corresponding formal spectral classes by

$$\begin{aligned}
J \in \widetilde{\text{Spec}}_{\mathbb{R}}(\mathfrak{A}) &\Leftrightarrow \frac{1}{z} \left(\left(\frac{1+z}{2} \right)^2 + \left(\frac{1-z}{2} \right)^2 J^2 \right) \text{ is invertible in } \mathfrak{A}[z, z^{-1}], \\
J \in \widetilde{\text{Evol}}(\mathfrak{A}) &\Leftrightarrow J^2 = -1, \\
Q \in \widetilde{\text{Spec}}_{i\mathbb{R}}(\mathfrak{A}) &\Leftrightarrow \frac{1+z}{2} + \frac{1-z}{2}Q = \Lambda(z, Q) \text{ is invertible in } \mathfrak{A}[z, z^{-1}], \\
Q \in \widetilde{\text{Spec}}_{\mathbb{C}^-}(\mathfrak{A}) &\Leftrightarrow \frac{1+z}{2} + \frac{1-z}{2}Q = \Lambda(z, Q) \text{ is invertible in } \mathfrak{A}[z], \\
Q \in \widetilde{\text{Invol}}(\mathfrak{A}) &\Leftrightarrow Q^2 = 1, \\
S \in \widetilde{\text{Spec}}_{\mathbb{R}}(\mathfrak{A}) &\Leftrightarrow \frac{1}{z} \left(\left(\frac{1+z}{2} \right)^2 - \left(\frac{1-z}{2} \right)^2 S \right) = \Lambda(z, S, z^{-1}) \text{ is invertible in } \mathfrak{A}[z, z^{-1}] \\
P \in \widetilde{\text{Spec}}_{\frac{1}{2}+i\mathbb{R}}(\mathfrak{A}) &\Leftrightarrow (1-P) + Pz \text{ is invertible in } \mathfrak{A}[z, z^{-1}], \\
P \in \widetilde{\text{Spec}}_{\frac{1}{2}-\mathbb{C}^-}(\mathfrak{A}) &\Leftrightarrow (1-P) + Pz \text{ is invertible in } \mathfrak{A}[z], \\
P \in \widetilde{\text{Idem}}(\mathfrak{A}) &\Leftrightarrow P^2 = P, \\
T \in \widetilde{\text{Spec}}_{\frac{1}{4}-\mathbb{R}}(\mathfrak{A}) &\Leftrightarrow 1 - (1-z)(1-z^{-1})T \text{ is invertible in } \mathfrak{A}[z, z^{-1}], \\
W \in \widetilde{\text{Spec}}_{\mathbb{C} \setminus \mathbb{D}^1}(\mathfrak{A}) &\Leftrightarrow 1 + zW \text{ is invertible in } \mathfrak{A}[z].
\end{aligned}$$

2.5. If $\mathfrak{A}$ is a locally convex algebra then the formal spectral classes and they ordinary counterparts are the same. Indeed, the continuity of the resolvent terms implies smoothness by the resolvent identity, hence the existence of the appropriate Fourier series, and, conversely, the existence of the expansions implies continuity.

Objective 2.6. We want to establish the spectral correspondences and decompositions as in point 2.2 for the formal spectral classes.

3. ELEMENTARY CALCULATIONS

A. Sign and square root.

Definition 3.1. For $Q \in \widetilde{\text{Spec}}_{i\mathbb{R}}(\mathfrak{A})$ we define

$$\text{sgn } Q = \int \frac{\frac{1-z}{2} + \frac{1+z}{2}Q}{\frac{1+z}{2} + \frac{1-z}{2}Q} \frac{|dz|}{2\pi} = \int \frac{\Lambda(-z, Q)}{\Lambda(z, Q)} \frac{|dz|}{2\pi}.$$

Proposition 3.2. If $Q \in \widetilde{\text{Spec}}_{i\mathbb{R}}(\mathfrak{A})$ then $-Q, Q^{-1} \in \widetilde{\text{Spec}}_{i\mathbb{R}}(\mathfrak{A})$. Q commutes with $\text{sgn } Q$. $\text{sgn } -Q = -\text{sgn } Q$ and $\text{sgn } Q^{-1} = \text{sgn } Q$. Moreover,

$$(\text{sgn } Q)^2 = 1.$$

Proof. Substituting $z = -1$ we see that Q^{-1} exists. From the identities $\Lambda(z, -Q) = z\Lambda(z^{-1}, Q)$ and $\Lambda(z, Q^{-1}) = Q^{-1}\Lambda(-z, Q)$ the first statement follows. Furthermore, Q and $\text{sgn } Q$ commute, because Q commutes with the integrand in $\text{sgn } Q$. The identities

$$\frac{\Lambda(-z, -Q)}{\Lambda(z, -Q)} \frac{|dz|}{2\pi} = -\frac{\Lambda(-z^{-1}, Q)}{\Lambda(z^{-1}, Q)} \frac{|d(z^{-1})|}{2\pi} \quad \text{and} \quad \frac{\Lambda(-z, Q^{-1})}{\Lambda(z, Q^{-1})} \frac{|dz|}{2\pi} = \frac{\Lambda(z, Q)}{\Lambda(-z, Q)} \frac{|d(-z)|}{2\pi}$$

integrated prove the first and second equalities, respectively.

The critical one is the involution property. We give several proofs.

“Matrix algebraic” proof. Let $H_{1/2} = \sum_{s \in \mathbb{Z} + \frac{1}{2}} (\text{sgn } s) \mathbf{e}_{s,s}$ be the $(\mathbb{Z} + \frac{1}{2}) \times (\mathbb{Z} + \frac{1}{2})$ matrix of the odd Hilbert transform. Let $U \left(\frac{1+Q}{2} z^{-1/2} + \frac{1-Q}{2} z^{1/2} \right) = \sum_{s \in \mathbb{Z} + \frac{1}{2}} \frac{1+Q}{2} \mathbf{e}_{s-\frac{1}{2},s} - \frac{1-Q}{2} \mathbf{e}_{s+\frac{1}{2},s}$ be the $\mathbb{Z} \times (\mathbb{Z} + \frac{1}{2})$ matrix of the action of multiplication by $\left(\frac{1+Q}{2} + \frac{1-Q}{2} z \right) z^{-1/2}$. According to our assumption this has a $(\mathbb{Z} + \frac{1}{2}) \times \mathbb{Z}$ inverse matrix representing the action of multiplication by $z^{1/2} \left(\frac{1+Q}{2} + \frac{1-Q}{2} z \right)^{-1}$. So we can consider the matrix

$$B \left(\frac{1+Q}{2} z^{-1/2} + \frac{1-Q}{2} z^{1/2} \right) = U \left(\frac{1+Q}{2} z^{-1/2} + \frac{1-Q}{2} z^{1/2} \right) H_{1/2} U \left(\frac{1+Q}{2} z^{-1/2} + \frac{1-Q}{2} z^{1/2} \right)^{-1}.$$

Due to the special shape of the matrices involved it is easy to see this is an involution which is the same as the even Hilbert transform $H = \sum_{s \in \mathbb{Z}} (\text{sgn } s) \mathbf{e}_{s,s}$ except in the 0th column. This special shape implies that the diagonal element in the 0th column is an involution. On the other hand, it is easy to see that this diagonal element is exactly $\text{sgn } Q$.

“Resolvent algebraic” proof. As

$$(6) \quad 1 - (\text{sgn } Q)^2 = \iint \left(1 - \frac{\Lambda(-z, Q)}{\Lambda(z, Q)} \frac{\Lambda(-w, Q)}{\Lambda(w, Q)} \right) \frac{|dz|}{2\pi} \frac{|dw|}{2\pi},$$

we should show that this integral is 0. This, however, follows from the key identity

$$(7) \quad 1 - \frac{\Lambda(-z, Q)}{\Lambda(z, Q)} \frac{\Lambda(-w, Q)}{\Lambda(w, Q)} = \left(\frac{z+w}{2} \right) \frac{1-Q^2}{\Lambda(z, Q)\Lambda(w, Q)} = \\ = \frac{1}{2} \left[\frac{z+w}{z-w} \right] \frac{(z-w)(1-Q^2)}{\Lambda(z, Q)\Lambda(w, Q)} = \frac{1}{2} \left[\frac{z+w}{z-w} \right] \left(\frac{(z-1)(1-Q^2)}{\Lambda(z, Q)} - \frac{(w-1)(1-Q^2)}{\Lambda(w, Q)} \right),$$

which does makes sense in $\mathfrak{A}[[z^{-1}, z]][[w^{-1}, w]]$. Indeed, (6) can be continued as

$$= \iint \left[\frac{z+w}{z-w} \right] \frac{(z-1)(1-Q^2)}{2\Lambda(z, Q)} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi} - \iint \left[\frac{z+w}{z-w} \right] \frac{(w-1)(1-Q^2)}{2\Lambda(w, Q)} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi}.$$

Executing the integrals we find

$$= \iint 0 \cdot \frac{(z-1)(1-Q^2)}{2\Lambda(z, Q)} \frac{|dz|}{2\pi} - \iint 0 \cdot \frac{(w-1)(1-Q^2)}{2\Lambda(w, Q)} \frac{|dw|}{2\pi} = 0 - 0 = 0.$$

This proof, like the previous one, relies heavily on the nature of Laurent series. Nevertheless the argument can be modified such that formally it makes sense in the analytical and the algebraic case as well.

“Resolvent analytic” proof. According to the discussion about transformation kernels, (6) can be continued as follows:

$$= \lim_{t \nearrow 1} \iint \tilde{\mathcal{R}}(t, wz^{-1}) \left(1 - \frac{\Lambda(-z, Q)}{\Lambda(z, Q)} \frac{\Lambda(-w, Q)}{\Lambda(w, Q)} \right) \frac{|dz|}{2\pi} \frac{|dw|}{2\pi}.$$

By simple arithmetic in the integrand, this yields

$$= \lim_{t \nearrow 1} \left(\iint \tilde{\mathcal{H}}(t, wz^{-1}) \frac{(z-1)(1-Q^2)}{2\Lambda(z, Q)} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi} - \iint \tilde{\mathcal{H}}(t, wz^{-1}) \frac{(w-1)(1-Q^2)}{2\Lambda(w, Q)} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi} \right).$$

Executing the integrals we find

$$= \lim_{t \nearrow 1} \left(\int 0 \cdot \frac{(z-1)(1-Q^2)}{\Lambda(z, Q)} \frac{|dz|}{2\pi} - \int 0 \cdot \frac{(w-1)(1-Q^2)}{\Lambda(w, Q)} \frac{|dw|}{2\pi} \right) = 0 + 0 = 0,$$

yielding, ultimately, the identity. We remark that in the analytic case it would actually be simpler to use the kernel $\mathcal{R}(t, wz^{-1})$. $\square$

Definition 3.3. If $S \in \widetilde{\mathfrak{Spec}}_{\mathbb{R}}(\mathfrak{A})$ then we define the inverse square root operation as

$$\sqrt{S} = \int \frac{zS}{\left(\frac{1+z}{2}\right)^2 - \left(\frac{1-z}{2}\right)^2} \frac{|dz|}{2\pi} = \int \frac{S}{\Lambda(z, S, z^{-1})} \frac{|dz|}{2\pi}.$$

Proposition 3.4. Suppose that $S \in \widetilde{\mathfrak{Spec}}_{\mathbb{R}}(\mathfrak{A})$. Then $S^{-1} \in \widetilde{\mathfrak{Spec}}_{\mathbb{R}}(\mathfrak{A})$. The elements $\sqrt{S}$ and S commute with each other. $\sqrt{S^{-1}} = \sqrt{S}^{-1}$. Furthermore,

$$(8) \quad (\sqrt{S})^2 = S.$$

Proof. Substituting $z = -1$ into the resolvent term we see that S^{-1} exists. The identity $\frac{1}{\Lambda(z, S^{-1}, z^{-1})} = \frac{S}{\Lambda(-z, S, (-z)^{-1})}$ shows that $S^{-1} \in \widetilde{\mathfrak{Spec}}_{\mathbb{R}}(\mathfrak{A})$. Integrated, it yields $S\sqrt{S^{-1}} = \sqrt{S}$. If the square-root identity (8) holds then this implies $\sqrt{S^{-1}} = \sqrt{S}^{-1}$. So, it remains to prove (8). As

$$(9) \quad (\sqrt{S})^2 - S = \iint \left(\frac{S}{\Lambda(z, S, z^{-1})} \frac{S}{\Lambda(w, S, w^{-1})} - S \right) \frac{|dz|}{2\pi} \frac{|dw|}{2\pi},$$

we have to show that this integral is 0. This follows using the key identities

$$(10) \quad \frac{S}{\Lambda(z, S, z^{-1})} \frac{S}{\Lambda(w, S, w^{-1})} - S = S \frac{S - \Lambda(z, S, z^{-1}, 1, w, S, w^{-1})}{2\Lambda(z, S, z^{-1})\Lambda(w, S, w^{-1})} + S \frac{S - \Lambda(z, S, z^{-1}, 1, w^{-1}, S, w)}{2\Lambda(z, S, z^{-1})\Lambda(w, S, w^{-1})};$$

$$(11) \quad \frac{S - \Lambda(z, S, z^{-1}, 1, w, S, w^{-1})}{2\Lambda(z, S, z^{-1})\Lambda(w, S, w^{-1})} = \frac{1}{2} \left[\frac{zw^{-1} + 1}{zw^{-1} - 1} \right] \left(\frac{\Lambda(-z, S, z^{-1})}{\Lambda(z, S, z^{-1})} - \frac{\Lambda(-w, S, w^{-1})}{\Lambda(w, S, w^{-1})} \right);$$

$$(12) \quad \frac{S - \Lambda(z, S, z^{-1}, 1, w^{-1}, S, w)}{2\Lambda(z, S, z^{-1})\Lambda(w, S, w^{-1})} = \frac{1}{2} \left[\frac{zw + 1}{zw - 1} \right] \left(\frac{\Lambda(-z, S, z^{-1})}{\Lambda(z, S, z^{-1})} - \frac{\Lambda(w, S, -w^{-1})}{\Lambda(w, S, w^{-1})} \right).$$

Indeed, after we decomposed the the integrand in (9) according to (10–12), we can show that either parts are 0 as we did in the case of previous proof. $\square$

Proposition 3.5. $Q \in \widetilde{\mathfrak{Spec}}_{\mathbb{R}}(\mathfrak{A})$ if and only if $Q^2 \in \widetilde{\mathfrak{Spec}}_{\mathbb{R}}(\mathfrak{A})$. In this case

$$\operatorname{sgn} Q = Q^{-1} \sqrt{Q^2}.$$

Proof. The first statement follows from the equality $\Lambda(z, Q^2, z^{-1}) = \Lambda(z, Q)\Lambda(z^{-1}, Q)$. The identity statement follows from

$$\begin{aligned} \operatorname{sgn} Q &= \frac{\operatorname{sgn} Q}{2} + \frac{\operatorname{sgn} Q}{2} = \frac{1}{2} \int \frac{\Lambda(-z, Q)}{\Lambda(z, Q)} \frac{|dz|}{2\pi} + \frac{1}{2} \int \frac{\Lambda(-z^{-1}, Q)}{\Lambda(z^{-1}, Q)} \frac{|d(z^{-1})|}{2\pi} \\ &= \int \frac{1}{2} \left(\frac{\Lambda(-z, Q)}{\Lambda(z, Q)} + \frac{\Lambda(-z^{-1}, Q)}{\Lambda(z^{-1}, Q)} \right) \frac{|dz|}{2\pi} = \int \frac{Q}{\Lambda(z, Q^2, z^{-1})} \frac{|dz|}{2\pi} = Q^{-1} \sqrt{Q^2}. \end{aligned} \quad \square$$

B. Finer analysis of the resolvent terms.

Definition 3.6. a.) We define $|Q|_r = \sqrt{Q^2} = Q \operatorname{sgn} Q$.

b.) If $F \in \mathfrak{A}$ is an involution then an element $A \in \mathfrak{A}$ can be written in matrix form

$$\begin{bmatrix} \frac{1-F}{2} A \frac{1-F}{2} & \frac{1-F}{2} A \frac{1+F}{2} \\ \frac{1+F}{2} A \frac{1-F}{2} & \frac{1+F}{2} A \frac{1+F}{2} \end{bmatrix}.$$

Suppose that $Q \in \widetilde{\text{Spec}}_{\mathbb{R}}(\mathfrak{A})$. In the decomposition of $\mathfrak{A}$ along the involution $\text{sgn } Q$ the various components are denoted according to

$$\text{sgn } Q = \begin{bmatrix} -1_{-\text{sgn } Q} & \\ & 1_{\text{sgn } Q} \end{bmatrix}, \quad Q = \begin{bmatrix} -Q^- & \\ & Q^+ \end{bmatrix}, \quad |Q|_r = \begin{bmatrix} Q^- & \\ & Q^+ \end{bmatrix}.$$

Proposition 3.7. $Q^\pm \in \widetilde{\text{Spec}}_{\mathbb{R}}(\mathfrak{A}1_{\pm \text{sgn } Q})$, and $\text{sgn } Q^\pm = 1_{\pm \text{sgn } Q}$. $\text{sgn } |Q|_r = 1$.

Proof. The decomposition of Q along $\text{sgn } Q$ allows us to consider Q separately in the direct sum components of $\mathfrak{A}$. In particular, the sign integral splits, too, and it necessarily yields $\text{sgn } Q^+ = 1_{\text{sgn } Q}$ and $\text{sgn } -Q^- = -1_{\text{sgn } -Q}$. From this the statement follows immediately. $\square$

Proposition 3.8. If $Q \in \widetilde{\text{Spec}}_{\mathbb{R}}(\mathfrak{A})$ then

a.)

$$\frac{1}{\frac{1+z}{2} + \frac{1-z}{2}Q} = \frac{1}{\Lambda(z, Q)}$$

is given by

$$\begin{aligned} \dots + \begin{bmatrix} \left(\frac{Q^- - 1}{Q^- + 1}\right)^2 \frac{2}{Q^- + 1} & \\ & 0 \end{bmatrix} z^{-3} + \begin{bmatrix} \left(\frac{Q^- - 1}{Q^- + 1}\right) \frac{2}{Q^- + 1} & \\ & 0 \end{bmatrix} z^{-2} + \begin{bmatrix} \frac{2}{Q^- + 1} & \\ & 0 \end{bmatrix} z^{-1} + \\ + \begin{bmatrix} 0 & \\ \frac{2}{Q^+ + 1} \end{bmatrix} 1 + \begin{bmatrix} 0 & \\ \frac{2}{Q^+ + 1} \left(\frac{Q^+ - 1}{Q^+ + 1}\right) \end{bmatrix} z + \begin{bmatrix} 0 & \\ \frac{2}{Q^+ + 1} \left(\frac{Q^+ - 1}{Q^+ + 1}\right)^2 \end{bmatrix} z^2 + \dots \end{aligned}$$

b.)

$$\frac{\frac{1-z}{2} + \frac{1+z}{2}Q}{\frac{1+z}{2} + \frac{1-z}{2}Q} = \frac{\Lambda(-z, Q)}{\Lambda(z, Q)}$$

is given by

$$\begin{aligned} \dots + \begin{bmatrix} -2 \left(\frac{Q^- - 1}{Q^- + 1}\right)^3 & \\ & 0 \end{bmatrix} z^{-3} + \begin{bmatrix} -2 \left(\frac{Q^- - 1}{Q^- + 1}\right)^2 & \\ & 0 \end{bmatrix} z^{-2} + \begin{bmatrix} -2 \left(\frac{Q^- - 1}{Q^- + 1}\right) & \\ & 0 \end{bmatrix} z^{-1} + \\ + \begin{bmatrix} -1_{-\text{sgn } Q} & \\ & 1_{\text{sgn } Q} \end{bmatrix} 1 + \begin{bmatrix} 0 & \\ 2 \left(\frac{Q^+ - 1}{Q^+ + 1}\right) \end{bmatrix} z + \begin{bmatrix} 0 & \\ 2 \left(\frac{Q^+ - 1}{Q^+ + 1}\right)^2 \end{bmatrix} z^2 + \begin{bmatrix} 0 & \\ 2 \left(\frac{Q^+ - 1}{Q^+ + 1}\right)^3 \end{bmatrix} z^3 + \dots \end{aligned}$$

c.) In particular,

$$\frac{1}{1 + |Q|_r} = \int \frac{1}{2} \frac{1+z}{\Lambda(z, Q)} \frac{|dz|}{2\pi}.$$

Proof. a.) It is enough to consider the Q^+ part of the decomposition, because the other part follows from changing z to z^{-1} . So, we can suppose that $Q = |Q|$ and $\text{sgn } Q = 1$.

We try to figure out the coefficients in the expansion $\frac{1}{\Lambda(z, Q)} = \sum_{n \in \mathbb{Z}} a_n z^n$. The equality $\text{sgn } Q = 1$ means that

$$(13) \quad 0 = \frac{1 - \text{sgn } Q}{2} = \frac{1}{2} \int \left(1 - \frac{\Lambda(-z, Q)}{\Lambda(z, Q)}\right) \frac{|dz|}{2\pi} = \int \frac{1 - Q}{2} \frac{z}{\Lambda(z, Q)} \frac{|dz|}{2\pi} = \frac{1 - Q}{2} a_{-1}.$$

The product of $\Lambda(z, Q) = \frac{1+Q}{2} + \frac{1-Q}{2}z$ and $\sum_{n \in \mathbb{Z}} a_n z^n$ gives 1, so

$$(14) \quad \frac{1-Q}{2} a_{n-1} + \frac{1+Q}{2} a_n = \delta_{0,n} 1.$$

From (13) and the case $n = 0$ in (14) we obtain that $a_0 = \frac{2}{1+Q}$. After that, from (14) we find $a_{n+1} = \frac{Q-1}{Q+1} a_n$, yielding the positive-numbered coefficients. Then $F(z) = \sum_{n \in \mathbb{N}} a_n z^n$ already inverts $\Lambda(z, Q)$, hence from the uniqueness of the inverse $a(z) = F(z)$.

b,c.) By simple algebra, they follow from part a. $\square$

Proposition 3.9. a.) If $S \in \widetilde{\text{Spec}}_{\mathbb{R}^-}(\mathfrak{A})$ then

$$\frac{z}{\left(\frac{1+z}{2}\right)^2 - \left(\frac{1-z}{2}\right)^2 S} = \frac{1}{\Lambda(z, S, z^{-1})}$$

yields the expansion

$$\sqrt{S}^{-1} 1 + \sqrt{S}^{-1} \left(\frac{\sqrt{S} - 1}{\sqrt{S} + 1} \right) (z + z^{-1}) + \sqrt{S}^{-1} \left(\frac{\sqrt{S} - 1}{\sqrt{S} + 1} \right)^2 (z^2 + z^{-2}) + \dots$$

b.) In particular

$$\frac{1}{\sqrt{S} + 1} = \int \frac{1}{2} \frac{1+z}{\Lambda(z, S, z^{-1})} \frac{|dz|}{2\pi}.$$

Proof. a.) Take $Q = \sqrt{S}$. From Proposition 3.5 it follows that $\text{sgn } Q = 1$. Applying the identity $\frac{1}{\Lambda(z, S, z^{-1})} = \frac{1}{\Lambda(z, Q)\Lambda(z^{-1}, Q)}$ and Proposition 3.8 it follows that in the expansion the coefficient of $z^{\pm n}$ ($n \geq 0$) is

$$\left(\frac{2}{Q+1} \right)^2 \sum_{m \in \mathbb{N}} \left(\frac{Q-1}{Q+1} \right)^{n+2m} = \left(\frac{2}{Q+1} \right)^2 \left(\frac{Q-1}{Q+1} \right)^n \left(1 - \left(\frac{Q-1}{Q+1} \right)^2 \right)^{-1},$$

which simplifies as above. The rapid decrease of $\left(\frac{Q-1}{Q+1} \right)^s$ makes our computations legal.

b.) It follows from part a. $\square$

Proposition 3.10. a.) $Q \in \widetilde{\text{Spec}}_{\mathbb{C}^-}(\mathfrak{A})$ if and only if $Q \in \widetilde{\text{Spec}}_{i\mathbb{R}^-}(\mathfrak{A})$ and $\text{sgn } Q = 1$.

b.) $W \in \widetilde{\text{Spec}}_{\mathbb{C} \setminus \mathbb{D}^1}(\mathfrak{A})$ if and only if W^n is rapidly decreasing.

c.) If $Q \in \widetilde{\text{Spec}}_{\mathbb{C}^-}(\mathfrak{A})$ then $\frac{1-Q}{1+Q} \in \widetilde{\text{Spec}}_{\mathbb{C} \setminus \mathbb{D}^1}(\mathfrak{A})$. Conversely, if $W \in \widetilde{\text{Spec}}_{\mathbb{C} \setminus \mathbb{D}^1}(\mathfrak{A})$ then $\frac{1-W}{1+W} \in \widetilde{\text{Spec}}_{\mathbb{C}^-}(\mathfrak{A})$. This establishes a bijection.

Proof. (a) follows from Proposition 3.8.a. (b) holds because in those cases $(1 - Wz)^{-1} = \sum_{n \in \mathbb{N}} W^n z^n$ must hold. (c) follows from $\Lambda\left(z, \frac{1-W}{1+W}\right) = \frac{1+zW}{1+W}$ and $1 + \frac{1-Q}{1+Q}z = \frac{2\Lambda(z, Q)}{1+Q}$. $\square$

C. Our objective.

3.11. Now, it is easy to see that the propositions proven above are sufficient establish all the spectral correspondences as in 2.6. We merely define $|J|_i = \sqrt{-J^2}$, $\text{pol } J = J| - J^2|_i^{-1}$ idem $P = \frac{1}{2} - \frac{1}{2} \text{sgn} \left(\frac{1}{2} - P \right)$, $|P|_{\mathcal{F}} = \frac{1}{2} - \left| \frac{1}{2} - P \right|_r$, and $\sqrt[{\mathcal{F}}]{T} = \frac{1}{2} - \sqrt{\frac{1}{4} - T}$. Hence our objective is established.

This is, however, not to say that everything is just like for locally convex algebras:

D. Comparison to the case of locally convex algebras.

Proposition 3.12. If $\mathfrak{A}$ is a locally convex algebra then the condition that $S \in \text{Spec}_{\mathbb{R}^-}(\mathfrak{A})$ is equivalent to the condition that the function

$$\frac{1+t}{2} + \frac{1-t}{2}S$$

has a continuous inverse on $[-1, 1]$. (This is the same thing as to say that the segment connecting 1 and S is continuously invertible.) The square root can be expressed as

$$\sqrt{S} = \int_{t \in [-1, 1]} \frac{S}{\frac{1+t}{2} + \frac{1-t}{2}S} \frac{dt}{\pi \sqrt{1-t^2}}.$$

Proof. It follows by change of variables using $t = \frac{z+z^{-1}}{2}$. $\square$

3.13. If $S \in \mathfrak{Spec}_{\mathbb{C}^-}(\mathfrak{A})$ then $\frac{1+t}{2} + \frac{1-t}{2}S$ ($t \in [-1, 1]$) is clearly invertible. As a rapidly decreasing power series in t , using $t = \frac{z+z^{-1}}{2}$ and considering the coefficients of z^k in $\left(\frac{z+z^{-1}}{2}\right)^n$, it follows that $S \in \mathfrak{Spec}_{\mathbb{R}^-}(\mathfrak{A})$. Hence the inclusion $\mathfrak{Spec}_{\mathbb{C}^-}(\mathfrak{A}) \subset \mathfrak{Spec}_{\mathbb{R}^-}(\mathfrak{A})$ is true. In the general context this cannot be done so because the boundedness of the elements $\frac{1}{2^n} \binom{n}{k}$ is not always clear. Similar comment applies for $\mathfrak{Spec}_{\frac{1}{2}+\mathbb{C}^-}(\mathfrak{A}) \subset \mathfrak{Spec}_{\frac{1}{2}+\mathbb{R}^-}(\mathfrak{A})$.

4. FORMAL HOMOTOPIES

4.1. One expects certain natural behaviour from the operations above. For example, one expects to have a homotopy from Q to $\text{sgn } Q$ inside $\widetilde{\mathfrak{Spec}}_{\mathbb{R}}(\mathfrak{A})$. In general algebras there are no continuous variables, but one can come up with homotopies using formal variables. Let us remind that an element $Q \in \widetilde{\mathfrak{Spec}}_{\mathbb{R}}(\mathfrak{A})$ can be decomposed to a commuting pair, $\text{sgn } Q$ and a perturbation of 1 which is $|Q|_r$. But we may also consider this as a decomposition to the commuting pair $\text{sgn } Q$ and a perturbation of 0 which is

$$\text{pert}_r Q = \frac{|Q|_r - 1}{|Q|_r + 1} = - \int \frac{\frac{1+z}{2} - \frac{1-z}{2}Q}{\frac{1+z}{2} + \frac{1-z}{2}Q} \frac{|dz|}{2\pi} = - \int \frac{\Lambda(z, -Q)}{\Lambda(z, Q)} \frac{|dz|}{2\pi}.$$

If we replace $\text{pert}_r Q$ by $t \text{pert}_r Q$ in the decomposition then we obtain a homotopy, this appears as $K(t, -1, Q)$ in what follows.

Definition 4.2. We define

$$\begin{aligned} K(t, z, Q) &= \frac{1 + \text{sgn } Q}{2} \frac{\Lambda(tz, |Q|_r)}{\Lambda(t, |Q|_r)} + \frac{1 - \text{sgn } Q}{2} z \frac{\Lambda(tz^{-1}, |Q|_r)}{\Lambda(t, |Q|_r)}, \\ H(t, z, Q) &= \frac{1 + \text{sgn } Q}{2} \frac{\Lambda(t, |Q|_r)}{\Lambda(tz, |Q|_r)} + \frac{1 - \text{sgn } Q}{2} z^{-1} \frac{\Lambda(t, |Q|_r)}{\Lambda(tz^{-1}, |Q|_r)}, \\ L(t, z, Q) &= \frac{1 + \text{sgn } Q}{2} \Lambda(tz, |Q|_r) + \frac{1 - \text{sgn } Q}{2} z \Lambda(tz^{-1}, |Q|_r), \\ G(t, z, Q) &= \frac{1 + \text{sgn } Q}{2} \frac{1}{\Lambda(tz, |Q|_r)} + \frac{1 - \text{sgn } Q}{2} z^{-1} \frac{1}{\Lambda(tz^{-1}, |Q|_r)}. \end{aligned}$$

Proposition 4.3. The expressions $K(t, z, Q)$ and $H(t, z, Q)$ are multiplicative inverses of each other.

$$K(t, z, Q) = \frac{\Lambda(z, \text{sgn } Q, t, |Q|_r)}{\Lambda(t, |Q|_r)} = \Lambda\left(z, \frac{\Lambda(-t, |Q|_r)}{\Lambda(t, |Q|_r)} \text{sgn } Q\right).$$

$$K(t, 1, Q) = 1, \quad K(1, -1, Q) = Q, \quad K(0, -1, Q) = \text{sgn } Q, \quad K(-1, -1, Q) = Q^{-1}.$$

$$H(t, z, Q) = \frac{\Lambda(t, |Q|_r) \Lambda(z^{-1}, \text{sgn } Q, t, |Q|_r)}{\Lambda(tz, |Q|_r) \Lambda(tz^{-1}, |Q|_r)}.$$

Similarly, the expressions $L(t, z, Q)$ and $G(t, z, Q)$ are inverses.

$$L(t, z, Q) = \Lambda(z, \text{sgn } Q, t, |Q|_r) = \Lambda(t, \text{sgn } Q, z, Q).$$

$$L(t, 1, Q) = \Lambda(t, |Q|_r), L(1, -1, Q) = Q, L(0, -1, Q) = \frac{1}{2}(Q + \operatorname{sgn} Q), L(-1, -1, Q) = \operatorname{sgn} Q.$$

$$G(t, z, Q) = \frac{\Lambda(z^{-1}, \operatorname{sgn} Q, t, |Q|_r)}{\Lambda(tz, |Q|_r) \Lambda(tz^{-1}, |Q|_r)}.$$

Proof. The computation is easy if we notice that in the defining formulas the coefficients of $\frac{1+\operatorname{sgn} Q}{2}$ and $\frac{1-\operatorname{sgn} Q}{2}$ live separate lives because $\operatorname{sgn} Q$ is an involution. $\square$

The properties of K show that $\frac{\Lambda(-t, |Q|_r)}{\Lambda(t, |Q|_r)} \operatorname{sgn} Q$ is homotopy from Q ($t = 1$) to $\operatorname{sgn} Q$ ($t = 0$) inside $\widetilde{\operatorname{Spec}}_{\mathbb{R}}(\mathfrak{A})$. Here the meaning of “inside” is that the whole expression satisfies the appropriate spectral condition.

Proposition 4.4.

$$\begin{aligned} K(t, w, Q) &= \int \mathcal{P}(t, z) \frac{\Lambda(zw, Q)}{\Lambda(z, Q)} \frac{|dz|}{2\pi}, \\ H(t, w, Q) &= \int \mathcal{P}(t, z) \frac{\Lambda(z, Q)}{\Lambda(zw, Q)} \frac{|dz|}{2\pi}, \\ G(t, w, Q) &= \int \mathcal{L}(t, z) \frac{1}{\Lambda(zw, Q)} \frac{|dz|}{2\pi}. \end{aligned}$$

Proof. This follows from the series expansion in Proposition 3.8. $\square$

Remark 4.5. In locally convex algebras the controllability of the powers of $\frac{z+z^{-1}}{2}$ makes possible to consider

$$\Lambda(z, \Lambda(t, |Q|_r) \operatorname{sgn} Q) = \frac{1 + \operatorname{sgn} Q}{2} \Lambda(\Lambda(z, t), |Q|_r) + \frac{1 - \operatorname{sgn} Q}{2} z \Lambda(\Lambda(z^{-1}, t), |Q|_r),$$

whose inverse turns out to be

$$\frac{\Lambda(z^{-1}, \Lambda(t, |Q|_r) \operatorname{sgn} Q)}{\Lambda(\Lambda(z, t), |Q|_r) \Lambda(\Lambda(z^{-1}, t), |Q|_r)}.$$

This shows that $\Lambda(t, |Q|_r) \operatorname{sgn} Q = \frac{1+t}{2} \operatorname{sgn} Q + \frac{1-t}{2} Q$ is also a formal homotopy between Q ($t = -1$) and $\operatorname{sgn} Q$ ($t = 1$) inside $\widetilde{\operatorname{Spec}}_{\mathbb{R}}(\mathfrak{A})$.

4.6. Similarly, we can contract elements inside $\widetilde{\operatorname{Spec}}_{\mathbb{R}}(\mathfrak{A})$ to 1. For $S \in \widetilde{\operatorname{Spec}}_{\mathbb{R}}(\mathfrak{A})$ consider

$$C(t, S) = \left(\frac{1 + t \frac{\sqrt{S}-1}{\sqrt{S}+1}}{1 - t \frac{\sqrt{S}-1}{\sqrt{S}+1}} \right)^2, \quad \sqrt{C(t, S)} = \frac{1 + t \frac{\sqrt{S}-1}{\sqrt{S}+1}}{1 - t \frac{\sqrt{S}-1}{\sqrt{S}+1}}.$$

The substitution $t \mapsto -t$ inverts them multiplicatively. In fact, the corresponding loops invert:

Proposition 4.7. For $S \in \widetilde{\operatorname{Spec}}_{\mathbb{R}}(\mathfrak{A})$

$$\frac{1}{\Lambda(w, C(t, S), w^{-1})} = \frac{\sqrt{S}}{\sqrt{C(t, S)}} \int \mathcal{P}(t, z) \frac{1}{\Lambda(zw, S, (zw)^{-1})} \frac{|dz|}{2\pi}.$$

Remark 4.8. In locally convex algebras alternative contracting paths are rather trivial to find. It is more interesting to see that the class of loops of type

$$\frac{1}{\frac{1}{z} \left(\left(\frac{1+z}{2} \right)^2 A^{-1} - \left(\frac{1-z}{2} \right)^2 B^{-1} \right)}$$

remains invariant to the Poisson kernel. For $t = 0$ they contract to the geometric mean

$$\sqrt{A \cdot B} = \int \frac{1}{\frac{1}{z} \left(\left(\frac{1+z}{2} \right)^2 A^{-1} - \left(\frac{1-z}{2} \right)^2 B^{-1} \right)} \frac{|dz|}{2\pi}.$$

5. CALCULATIONS WITHOUT $\frac{1}{2}$

As we have seen much can be generalized to the case $\frac{1}{2} \in \mathfrak{A}$. It is natural to ask that what happens in the lack of $\frac{1}{2}$. Then only the lower portion of (5) can be generalized. Again, the idempotent and the $\mathcal{F}$ -square-root identities are key properties.

Definition 5.1. For $P \in \mathfrak{Spec}_{\frac{1}{2} + i\mathbb{R}}(\mathfrak{A})$ we define

$$\text{idem } P = \int \frac{Pz}{1 - P + Pz} \frac{|dz|}{2\pi}.$$

Proposition 5.2. If $P \in \mathfrak{Spec}_{\frac{1}{2} + i\mathbb{R}}(\mathfrak{A})$ then $1 - 2P$ is invertible, $1 - P, \frac{-P}{1 - 2P} \in \mathfrak{Spec}_{\frac{1}{2} + i\mathbb{R}}(\mathfrak{A})$, and $\text{idem}(1 - P) = 1 - \text{idem } P$, $\text{idem} \frac{-P}{1 - 2P} = \text{idem } P$. Furthermore,

$$(\text{idem } P)^2 = \text{idem } P.$$

Proof. The invertibility statement follows from the substitution $z = -1$. The identities $P + (1 - P)z = (P + (1 - P)z^{-1})z$ and $(1 - \frac{-P}{1 - 2P}) + \frac{-P}{1 - 2P}z = (1 - 2P)^{-1}(1 - P + P(-z))$ imply the spectral statements. The identities

$$\begin{aligned} \frac{Pz}{(1 - P) + Pz} \frac{|dz|}{2\pi} &= \left(1 - \frac{(1 - P)z^{-1}}{P + (1 - P)z^{-1}} \right) \frac{|d(z^{-1})|}{2\pi} \\ \frac{\frac{-P}{1 - 2P}z}{(1 - \frac{-P}{1 - 2P}) + \frac{-P}{1 - 2P}z} \frac{|dz|}{2\pi} &= \frac{P(-z)}{1 - P + P(-z)} \frac{|d(-z)|}{2\pi} \end{aligned}$$

integrated prove the first and second equalities. We can prove the idempotent identity in several ways:

Matrix algebraic proof. We can proceed as before but have to conjugate not the Hilbert transform involution but the idempotent $\sum_{s \in -\mathbb{N} - \frac{1}{2}} \mathbf{e}_{s,s}$.

A direct algebraic proof. See Karoubi [2], Lemma III.1.23–24.

A resolvent algebraic proof. We should prove that

$$(15) \quad \text{idem } P(1 - \text{idem } P) = \iint \frac{Pz}{1 - P + Pz} \frac{1 - P}{1 - P + Pw} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi}$$

is equal to 0. It is natural try the proof along the steps

$$\begin{aligned} (16) \quad \frac{Pz}{1 - P + Pz} \frac{1 - P}{1 - P + Pw} &\sim \left[\frac{z + w}{2} \right] \frac{P(1 - P)}{(1 - P + Pz)(1 - P + Pw)} = \\ &= \frac{1}{2} \left[\frac{z + w}{z - w} \right] \left(\frac{(z - 1)P(1 - P)}{1 - P + Pz} - \frac{(w - 1)P(1 - P)}{1 - P + Pw} \right) \sim 0 \end{aligned}$$

except it seems to be plagued by $1/2$'s as before. We have to demonstrate that the use of division by 2 is of superficial nature in the proof. This can be done as follows.

For a Laurent series $a(z, w)$ we define $:a(z, w):_{z,w}$ by linear extension from

$$:z^n w^m:_{z,w} = z^{\max(n,m)} w^{\min(n,m)}.$$

Lemma 5.3. For any Laurent series $a(z, w)$

$$\iint a(z, w) \frac{|dz|}{2\pi} \frac{|dw|}{2\pi} = \iint :a(z, w):_{z,w} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi}.$$

For a symmetric Laurent series $a(z, w)$, ie. such that $b(z, w) = b(w, z)$, we define formally

$$: \left[\frac{z+w}{2} \right] a(z, w) :_{z,w} =: za(z, w) :_{z,w}.$$

Suppose that $a(z, w)$ is anti-symmetric in its variables, included that the coefficient of $z^n w^n$ is always 0. We define $: \frac{1}{2} \left[\frac{z+w}{z-w} \right] a(z, w) :_{z,w}$ formally to be as it should be according to our natural expectations. We have to check that the resulting expression is integral in terms of the coefficients. In the present case the definition yields by linear extension from

$$: \frac{1}{2} \left[\frac{z+w}{z-w} \right] (z^n w^m - z^m w^n) :_{z,w} = z^n w^m + 2 \sum_{0 < k < \frac{n-m}{2}} z^{n-k} w^{m+k} + \delta_{\frac{n+m}{2} \in \mathbb{Z}} z^{\frac{n+m}{2}} w^{\frac{n+m}{2}}$$

where $n > m$. By extension from good basis elements it easy to see the following lemmas:

Lemma 5.4. *For any symmetric Laurent series $a(z, w)$*

$$: \left[\frac{z+w}{2} \right] a(z, w) :_{z,w} =: \frac{1}{2} \left[\frac{z+w}{z-w} \right] (z-w)a(z, w) :_{z,w}.$$

Lemma 5.5. *For any Laurent series $a(z)$*

$$\iint : \frac{1}{2} \left[\frac{z+w}{z-w} \right] (a(z) - a(w)) :_{z,w} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi} = 0.$$

Now, it is easy to carry out the proof. From (15) we should pass to the “normal ordered” form, after which the subsequent manipulations as in (16) leading to 0 make sense. $\square$

Definition 5.6. For $T \in \mathfrak{Spec}_{\frac{1}{4}-\mathbb{R}^-}(\mathfrak{A})$ we define

$$\sqrt[4]{T} = \int \frac{(1+z)T}{1+(z-2+z^{-1})T} \frac{|dz|}{2\pi}.$$

Definition 5.7. Let $\mathfrak{A}\langle z \rangle^+$ be the space of formal Laurent series $a(z) = a_0 + \sum_{k=1}^{\infty} a_k \frac{z^k + z^{-k}}{2}$, and let $\mathfrak{A}[z]^+$ be the space of formal Laurent series $b(z) = b_0 + \sum_{k=1}^{\infty} b_k (z^k + z^{-k})$. Similarly, let $\mathfrak{A}\langle z \rangle^-$ be the space of formal Laurent series $a(z) = \sum_{k=1}^{\infty} a_{-k} \frac{z^k - z^{-k}}{2}$, and let $\mathfrak{A}[z]^-$ be the space of formal Laurent series $b(z) = \sum_{k=1}^{\infty} b_{-k} (z^k - z^{-k})$. It is easy to see that $\mathfrak{A}\langle z \rangle = \mathfrak{A}\langle z \rangle^+ \oplus \mathfrak{A}\langle z \rangle^-$ is a natural $\mathfrak{A}[z]^{\pm} = \mathfrak{A}[z]^+ \oplus \mathfrak{A}[z]^-$ -module, in fact, this action is $\mathbb{Z}_2$ -graded. Multiplication of $1 \in \mathfrak{A}\langle z \rangle$ yields a natural map from $\mathfrak{A}[z]^{\pm}$ into $\mathfrak{A}\langle z \rangle$:

$$1 \cdot \left(\sum_{k=1}^{\infty} b_{-k} (z^k - z^{-k}) + b_0 + \sum_{k=1}^{\infty} b_k (z^k + z^{-k}) \right) = \sum_{k=1}^{\infty} 2b_{-k} \frac{z^k - z^{-k}}{2} + b_0 + \sum_{k=1}^{\infty} 2b_k \frac{z^k + z^{-k}}{2}.$$

In fact, this notation can be extended to $b(z) \in \mathfrak{A}[z, z^{-1}]$ compatibly, by $1 \cdot z = \frac{z+z^{-1}}{2} + \frac{z-z^{-1}}{2}$, etc. Integration can be defined for elements of $\mathfrak{A}\langle z \rangle$ or $\mathfrak{A}[z]^{\pm}$, again it singles out 0th coefficient. We see that if $a(z) \in \mathfrak{A}\langle z \rangle$ and $b(z) \in \mathfrak{A}[z]^{\pm}$ as above then

$$\int a(z) \cdot b(z) \frac{|dz|}{2\pi} = \sum_{n \in \mathbb{Z}} a_n b_n.$$

For example, $\int \left(1 + \frac{z+z^{-1}}{2} \right) \cdot b(z) \frac{|dz|}{2\pi} = b_0 + b_1$. Furthermore, $\int b(z) \frac{|dz|}{2\pi} = \int 1 \cdot b(z) \frac{|dz|}{2\pi}$.

We can extend this formalism to multiple variables. The spaces $\mathfrak{A}\langle z \rangle \langle w \rangle$ and $\mathfrak{A}[z]^{\pm} [w]^{\pm}$ can be considered. In case of $\mathfrak{A}\langle z \rangle \langle w \rangle$ the colloquial notation like

$$\frac{zw^{-1} + wz^{-1}}{2} \equiv \frac{z+z^{-1}}{2} \frac{w+w^{-1}}{2} - \frac{z-z^{-1}}{2} \frac{w-w^{-1}}{2}$$

is allowed. However, an other space between $\mathfrak{A}\langle z \rangle \langle w \rangle$ and $\mathfrak{A}[z]^\pm[w]^\pm$ can be considered. Indeed, let $\mathfrak{A}\{z, w\}$ be the space of the formal combinations of the basis elements

$$1, \frac{z^n+z^{-n}}{2}, \frac{z^n-z^{-n}}{2}, \frac{w^n+w^{-n}}{2}, \frac{w^n-w^{-n}}{2}, \\ \frac{(z^n+z^{-n})(w^n+w^{-n})}{2}, \frac{(z^n+z^{-n})(w^n-w^{-n})}{2}, \frac{(z^n-z^{-n})(w^n+w^{-n})}{2}, \frac{(z^n-z^{-n})(w^n-w^{-n})}{2}$$

where $n, m \geq 1$. In this case colloquial notation like

$$zw^{-1} - wz^{-1} \equiv \frac{(z-z^{-1})(w+w^{-1})}{2} - \frac{(z+z^{-1})(w-w^{-1})}{2},$$

is allowed. There are natural $\mathfrak{A}[z]^\pm[w]^\pm$ -module homomorphisms $\mathfrak{A}[z]^\pm[w]^\pm \rightarrow \mathfrak{A}\{z, w\} \rightarrow \mathfrak{A}\langle z \rangle \langle w \rangle$ respecting grading.

5.8. The advantage of the terminology above is that it allows us to rewrite the definition of the $\mathcal{F}$ -square-root as the “manifestly real” expression

$$\sqrt[\mathcal{F}]{T} = \int \frac{(1 + \frac{z+z^{-1}}{2})T}{1 + (z - 2 + z^{-1})T} \frac{|dz|}{2\pi}.$$

Proposition 5.9. *If $T \in \mathfrak{Spec}_{\frac{1}{4}-\mathbb{R}}(\mathfrak{A})$ then $\frac{-T}{1-4T}$ is invertible and $\sqrt[\mathcal{F}]{\frac{-T}{1-4T}} = \frac{-\sqrt[\mathcal{F}]{T}}{1-2\sqrt[\mathcal{F}]{T}}$. Furthermore,*

$$(17) \quad \sqrt[\mathcal{F}]{T}(1 - \sqrt[\mathcal{F}]{T}) = T.$$

Proof. Substituting $z = -1$ into the resolvent term we see that $(1 - 4T)^{-1}$ exists. The identity $1 + (z - 2 + z^{-1})\frac{-T}{1-4T} = (1 - 4T)^{-1}(1 + ((-z) - 2 + (-z)^{-1})T)$ shows that $\frac{-T}{1-4T} \in \mathfrak{Spec}_{\frac{1}{4}-\mathbb{R}}(\mathfrak{A})$. If (17) holds then $(1 - 2\sqrt[\mathcal{F}]{T})^2 = 1 - 4T$, and it is sufficient to prove that $\sqrt[\mathcal{F}]{\frac{-T}{1-4T}} = \frac{\sqrt[\mathcal{F}]{T}-2T}{1-4T}$. This, however, follows from the identity

$$\frac{(1 + \frac{z+z^{-1}}{2})\frac{-T}{1-4T}}{1 + (z - 2 + z^{-1})\frac{-T}{1-4T}} = \frac{\frac{(1 + \frac{(-z)+(-z)^{-1}}{2})T}{1 + ((-z) - 2 + (-z)^{-1})T} - 2T}{1 - 4T}$$

integrated. So we have show the $\mathcal{F}$ -square-root identity (17). Now, as

$$(18) \quad \sqrt[\mathcal{F}]{T}(1 - \sqrt[\mathcal{F}]{T}) - T = \iint \frac{(1 + \frac{z+z^{-1}}{2})T}{1 + (z - 2 + z^{-1})T} \left(1 - \frac{(1 + \frac{w+w^{-1}}{2})T}{1 + (w - 2 + w^{-1})T} \right) - T \frac{|dz|}{2\pi} \frac{|dw|}{2\pi},$$

we should show that this latter term is 0. It would be natural to proceed along the steps

$$(19) \quad \frac{(1 + \frac{z+z^{-1}}{2})T}{1 + (z - 2 + z^{-1})T} \left(1 - \frac{(1 + \frac{w+w^{-1}}{2})T}{1 + (w - 2 + w^{-1})T} \right) - T = \\ = \frac{\left(\frac{z+z^{-1}}{2} - \frac{z+z^{-1}}{2}T - \frac{w+w^{-1}}{2}T + T + \frac{z+z^{-1}}{2} \frac{w+w^{-1}}{2}T \right) T(1 - 4T)}{(1 + (z - 2 + z^{-1})T)(1 + (w - 2 + w^{-1})T)} \\ \sim \frac{\left(\left[\frac{\frac{z+z^{-1}}{2} + \frac{w+w^{-1}}{2}}{2} \right] (1 - 2T) + \left(1 + \frac{z^{-1}w + zw^{-1}}{2} \right) T \right) T(1 - 4T)}{(1 + (z - 2 + z^{-1})T)(1 + (w - 2 + w^{-1})T)} \\ = \frac{1}{2} \left[\frac{z+w}{z-w} \right] \frac{\left(\left(\frac{z-z^{-1}}{2} - \frac{w-w^{-1}}{2} \right) (1 - 2T) + (zw^{-1} - z^{-1}w)T \right) T(1 - 4T)}{(1 + (z - 2 + z^{-1})T)(1 + (w - 2 + w^{-1})T)} \\ = \frac{1}{2} \left[\frac{z+w}{z-w} \right] \left(\frac{\frac{z-z^{-1}}{2}}{1 + (z - 2 + z^{-1})T} - \frac{\frac{w-w^{-1}}{2}}{1 + (w - 2 + w^{-1})T} \right) T(1 - 4T) \sim 0,$$

except we have to demonstrate that the use of $\frac{1}{2}$ is superficial.

A reordering operation is to be defined according to

$$:: z^n w^m ::_{z,w} = \frac{z^{\max(|n|,|m|)} + z^{-\max(|n|,|m|)}}{2} \frac{w^{\min(|n|,|m|)} + w^{-\min(|n|,|m|)}}{2}.$$

It applies to our standard $\mathfrak{A}[z]^\pm[w]^\pm$ -modules. In the context of $\mathfrak{A}\langle z \rangle \langle w \rangle$ it leaves only the $(++)$ -graded parts and reorders them..

Lemma 5.10. *a.) For any Laurent series $a(z, w) \in \mathfrak{A}\langle z \rangle \langle w \rangle$*

$$\iint a(z, w) \frac{|dz|}{2\pi} \frac{|dw|}{2\pi} = \iint :: a(z, w) ::_{z,w} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi}.$$

b.) If $:: a_1(z, w) ::_{z,w} = :: a_2(z, w) ::_{z,w}$ and $b(z, w) \in \mathfrak{A}[z]^+[w]^+$ is symmetric, ie. $b(z, w) = b(w, z)$, then $:: a_1(z, w) \cdot b(z, w) ::_{z,w} = :: a_2(z, w) \cdot b(z, w) ::_{z,w}$.

For any symmetric Laurent series $b(z, w) \in \mathfrak{A}[z]^+[w]^+$, ie. such that $b(z, w) = b(w, z)$, we define formally

$$:: \left[\frac{z+z^{-1}}{2} + \frac{w+w^{-1}}{2} \right] b(z, w) ::_{z,w} = :: \frac{z+z^{-1}}{2} b(z, w) ::_{z,w}.$$

Let us consider a Laurent series $c(z, w) \in (\mathfrak{A}\{z, w\})^-$ such that it is antisymmetric, ie. $c(z, w) = -c(w, z)$. Then $c(z, w)$ is a formal linear combination of the basis elements

$$c_{n,0} = \frac{z^n - z^{-n}}{2} - \frac{w^n - w^{-n}}{2} \quad \text{and} \quad c_{n,m} = \frac{(z^n - z^{-n})(w^m + w^{-m})}{2} - \frac{(w^n - w^{-n})(z^m + z^{-m})}{2}$$

where $n, m \geq 1$. For such $c(z, w)$ we can formally define $:: \frac{1}{2} \left[\frac{z+w}{z-w} \right] c(z, w) ::_{z,w}$ according to our natural expectations. Again, we have to check that the result is integral in terms of coefficients of $c(z, w)$. We just give some samples in the table

$:: \frac{1}{2} \left[\frac{z+w}{z-w} \right] c_{n,m} ::_{z,w}$	$m = 0$	$m = 1$	$m = 2$	$m = 3$
$n = 1$	d_0^1	$d_1^1 + d_0^0$	$2d_0^1$	$-d_2^2 + d_1^1 + 2d_0^2$
$n = 2$	$d_0^2 + d_1^1$	$2d_1^2 + 2d_0^1$	$d_2^2 + 2d_1^1 + d_0^0$	$2d_1^2 + 2d_0^1$
$n = 3$	$d_0^3 + 2d_1^2$	$2d_1^3 + d_2^2 + d_1^1 + 2d_0^2$	$2d_2^3 + 2d_1^2 + 2d_0^1$	$d_3^3 + 2d_2^2 + 2d_1^1 + d_0^0$

where $d_m^n = :: z^n w^m ::_{z,w}$. In fact, at first sight it looks more natural to choose $c(z, w)$ from the antisymmetric elements of $(\mathfrak{A}\langle z \rangle \langle w \rangle)^- = \mathfrak{A}\langle z \rangle^+ \langle w \rangle^- \oplus \mathfrak{A}\langle z \rangle^- \langle w \rangle^+$ but it turns out that the coefficients in $:: \frac{1}{2} \left[\frac{z+w}{z-w} \right] c(z, w) ::_{z,w}$ would fail to be integral.

Lemma 5.11. *For any symmetric Laurent series $b(z, w) \in \mathfrak{A}[z]^+[w]^+$*

$$:: \left[\frac{z+z^{-1}}{2} + \frac{w+w^{-1}}{2} \right] b(z, w) ::_{z,w} = :: \frac{1}{2} \left[\frac{z+w}{z-w} \right] \left(\frac{z-z^{-1}}{2} - \frac{w-w^{-1}}{2} \right) \cdot b(z, w) ::_{z,w},$$

and

$$:: \left(1 + \frac{zw^{-1} + wz^{-1}}{2} \right) b(z, w) ::_{z,w} = :: \frac{1}{2} \left[\frac{z+w}{z-w} \right] (zw^{-1} - wz^{-1}) \cdot b(z, w) ::_{z,w}.$$

Lemma 5.12. *For any Laurent series $a(z) \in \mathfrak{A}\langle z \rangle^-$*

$$\iint :: \frac{1}{2} \left[\frac{z+w}{z-w} \right] (a(z) - a(w)) ::_{z,w} \frac{|dz|}{2\pi} \frac{|dw|}{2\pi} = 0.$$

Now, it is easy to carry out the proof. From (18) we should pass to the “normal ordered” form, after which the subsequent manipulations leading to 0 make sense. $\square$

Proposition 5.13. $P \in \mathfrak{Spec}_{\frac{1}{2} + i\mathbb{R}}(\mathfrak{A})$ if and only if $P(1-P) \in \mathfrak{Spec}_{\frac{1}{4} - \mathbb{R}^-}(\mathfrak{A})$. Furthermore,

$$\sqrt[\mathcal{F}]{P(1-P)} = P + \text{idem } P - 2P \text{idem } P.$$

Consequently,

$$\text{idem } P = \frac{\sqrt[\mathcal{F}]{P(1-P)} - P}{1 - 2P}.$$

Proof. The decomposition $1 + (z - 2 + z^{-1})P(1-P) = (1 - P + Pz)(1 - P + Pz^{-1})$ implies the spectral statement. The equality follows from the identity

$$\frac{\left(1 + \frac{z+z^{-1}}{2}\right)P(1-P)}{1 + (z - 2 + z^{-1})P(1-P)} + \frac{\frac{z-z^{-1}}{2}P(1-P)(1-2P)}{1 + (z - 2 + z^{-1})P(1-P)} = P + \frac{Pz}{1 - P + Pz}(1 - 2P)$$

integrated. $\square$

Then we let $|P|_{\mathcal{F}} = \sqrt[\mathcal{F}]{P(1-P)}$. Further statements can be proven parallel to the case with $\frac{1}{2}$, except the formulas are less customary. E. g., the analogy of Proposition 3.9 is

Proposition 5.14. For $T \in \mathfrak{Spec}_{\frac{1}{4} - \mathbb{R}^-}(\mathfrak{A})$

$$\frac{1}{1 + (z - 2 + z^{-1})T} = \frac{1}{1 - 2\sqrt[\mathcal{F}]{T}} \left(1 + \frac{-\sqrt[\mathcal{F}]{T}}{1 - \sqrt[\mathcal{F}]{T}}(z + z^{-1}) + \left(\frac{-\sqrt[\mathcal{F}]{T}}{1 - \sqrt[\mathcal{F}]{T}} \right)^2 (z^2 + z^{-2}) + \dots \right)$$

and

$$\frac{1}{1 - \sqrt[\mathcal{F}]{T}} = \int \frac{1 + z}{1 + (z - 2 + z^{-1})T} \frac{|dz|}{2\pi}.$$

REFERENCES

- [1] M. Haase: *The Functional Calculus for Sectorial Operators*. Operator Theory: Advances and Applications 169. Birkhäuser Verlag, Basel, 2006.
- [2] M. Karoubi: *K-theory. An introduction*. Grundlehren der mathematischen Wissenschaften, 226. Springer Verlag, Berlin, Heidelberg, New York, 1978.
- [3] C. Martínez Carracedo, M. Sanz Alix: *The theory of fractional powers of operators*. North-Holland Mathematics Studies, 187. North-Holland, Amsterdam, 2001.

DEPARTMENT OF GEOMETRY, EÖTVÖS UNIVERSITY, PÁZMÁNY PÉTER S. 1/C, BUDAPEST, H-1117, HUNGARY

E-mail address: lakos@cs.elte.hu